

PROTECTIVE EFFECT OF *AGERATUM CONYZOIDES* ON CARBON TETRACHLORIDE – INDUCED HEPATIC DAMAGE IN RATS

Kiran Kushwaha^{*1}, Varun Jain^{*2}

^{*1,2}SAM Global University Raisen, India.

ABSTRACT

The present study was conducted to examine the Hepatoprotective activity against the carbon tetrachloride toxicity in Wister rats. The Petroleum ether, ethyl acetate and methanol extract of *Ageratum canyzoides* was prepared and evaluated for phytochemical screening. The serum level of glutamic oxaloacetate transaminase (SGOT) Glutamicpyruvate transaminase (SGPT) and bilirubin were investigated for the assessment of hepatoprotective activity of ethyl acetate of extract *Ageratumconyzoides*.

Methods

The whole plant *Ageratum conyzoides* was collected from the local surroundings of Bhopal region, during the month of August - October 2020. The plant was identified by Dr. Jagrati Tripathi - Professor of Botany, Govt. College Khemlasha Sagar. The voucher specimens

Results

Additionally, the histological changes in liver were studied. The primary phytochemical investigation of the extract of *Ageratumconyzoides* revealed the presence of Flavonoids, tannins, and carbohydrates. Pretreatment with ethyl acetate extract of *Ageratumconyzoides* showed a significant ($P < 0.05$) decrease in serum SGOT, SGPT and bilirubin when compared to control group rats treated with CCl_4 in dose dependent manner. The outcomes of histology study revealed that there was significant reversal of histological functions of liver. In conclusion, the findings of study validated that the *Ageratum canyzoides* can improve CCl_4 -induced hepatic toxicity.¹⁻⁴

Keywords: *Ageratum Conyzoides*, SGOT, SGPT, Glutamic Oxaloacetate, Hepatotoxins Hepatoprotective Flavonoids.

I. INTRODUCTION

Liver regulates various important metabolic functions and damage is associated with distortion of their metabolic function. Today liver damage is a very common ailment in the world resulting in serious disabilities ranging from severe metabolic disorders to even mortality.

Antioxidants such as GSH, ascorbic acid, and Vitamin E and enzymes such as SOD catalase and GPX acts against oxidative process. Hepatocytes which make most of the liver structure are very active in the metabolism of exogenous chemicals and is one of the major reasons. Carbon tetra methyl chloride is one of the most commonly used hepatotoxins in the experimental study of liver diseases.⁵⁻⁸

II. MATERIAL AND METHODS

Preparation of Plant Material

The whole plant *Ageratum conyzoides* was collected from the local surroundings of Bhopal region, during the month of August - October 2020. The plant was identified by Dr. Jagrati Tripathi - Professor of Botany, Govt. College Khemlasha Sagar. The voucher specimens are kept in the P.G. Department of Zoology, Career College, Bhopal. The whole plant was washed thoroughly under running tap water, the dried plant Material powdered and subjected to extraction.

Preparation of Extract: -

The extract was done by maceration using petroleum ether, ethyl acetate and methanol. The extract obtained was evaporated in rotary evaporator to get a powder. The extract then subjected to phytochemical analysis to detect the chemical constitution present in each extract.

Animals: -

Male wister rats (135-108g) were used for evaluation of hepatoprotective activity. The animals were housed in polypropylene cages at $25^\circ\text{C} \pm 1^\circ\text{C}$ with the relative humidity of $55 \pm 5\%$ under 12h/12h light /dark cycle. They

received a standard chow, water and labium throughout the experiments, according to the International ethical guidelines for the care of laboratory animals.¹⁰⁻¹². The study protocol was approved by the Institutional Animal Ethical committee.

CCl₄ induced hepatotoxicity-Assessment of hepatoprotective activity was carried on albino rats. The animals were segregated in four groups consisting of six animals each.

Group I served as normal control receiving 5% CMC (10 ml / kg)

Group II received (1ml/ kg ip) equal volume of olive oil(50%v/v) for two successive days and were maintained as CCl₄ group.

Group III animals were treated orally for seven days with suspension of ethyl acetate extract (100mg/kg)

Group IV animals were treated with CCl₄ and ethyl acetate extract (100mh/kg).

After the treatment the blood samples were collected via retro orbital and the serum was separated by centrifugation at 2500 rpm for 15 min for the estimation of biochemical enzyme. After collecting the blood, the liver was separated and weighted (20-25g).¹³

Biochemical determination: -

Biochemical Parameters such as serum glutamate oxaloacetate transaminase (SGOT) serum glutamate pyruvate transaminase (SGPT) and serum bilirubin were determined.

Histopathology: - Liver was excised quickly and fixed in 10% buffered neutral formalin and processed for paraffin embedding following the standard micro techniques. Section of liver is stained with alumhaematoxylin and eosin were observed microscopically for histopathology changes.

Statistical analysis: -The results are expressed as mean \pm SEM of six animals from each group. The data was evaluated by one way ANOVA followed by Turkeys multiple Comparison tests. P values <0.05 were considered statistically significant.

Results and Discussion: -

Phytochemicals Screening of *Ageratum conyzoides*: -

Presence of secondary metabolites might be a useful indicator of both efficacy and potential toxicity: hence test for the presence of phytochemical classes with known bioactivity was done with preliminary phytochemical investigation of the extracts of whole plants. The treatment with ethyl acetate extracts of *Ageratum conyzoides* at dose (100mg/kg) has decreased. Thus, the activity of SGOT, SGPT and total bilirubin in CCl₄ treated rats was examined.

Table 1: Effect of methanolic extract of *A.conyzoides* isolated compound on biochemical parameters of liver in rats against CCl₄ administration.

S. No.	Treatment	Dose	SGOT/AST(IU/L)	SGPT/ALT (IU/L)	ALP (IU/L)	Bilirubin (IU/L)
1	Vehicle Saline	10 ml/kg	32.1 \pm 4.845	28.8 \pm 3.746	105.4 \pm 8.016	0.76 \pm 0.048
2	Control (CCl ₄)	-	136.6 \pm 5.680*	132.6 \pm 2.581*	302.3 \pm 9.872*	1.81 \pm 0.111*
3	Silymarine	100 mg/kg	44.1 \pm 3.920**	38.5 \pm 8.336**	125.1 \pm 7.833**	0.80 \pm 0.054**
4	MEEA	200 mg/kg	84.6 \pm 4.226**	78.6 \pm 6.377**	223.5 \pm 9.027**	1.08 \pm 0.057**
5	MEEA	400 mg/kg	69.1 \pm 4.792**	63.1 \pm 6.177**	161.8 \pm 5.344**	0.93 \pm 0.031**

Each group consists of six animals (N=6).

MEEA = Methanolic extract of *A.conyzoides**P<0.001 as compared to vehicle treated group

** P<0.001 as compared to CCl₄ treated group

P<0.001 as compared to vehicle treated group

Table 2: Effect of isolated compound on biochemical parameters of liver in rats against CCl₄ administration.

S. No.	Treatment	Dose	SGOT/AST(IU/L)	SGPT/ALT(IU/L)	ALP (IU/L)	BILIRUBIN (IU/L)
1	VehicleSaline	10 ml/kg	32.1±4.845	28.8±3.746	105.4±8.016	0.76±0.048
2	Control (CCl ₄)	-	136.6±5.680*	132.6±2.581*	302.3±9.872*	1.81±0.111*
3	Silymarine	100 mg/kg	44.1±3.920**	38.5±8.336**	125.1±7.833**	0.80±0.054**
4	Isolated fraction	100 mg/kg	75.36±4.313**	61.76±5.243**	197.5±7.152**	0.902±0.042**
5	Isolated fraction	200 mg/kg	56.1±3.654**	52.4±4.231**	141.1±3.253**	0.877±0.021**

III. DISCUSSION

According to Rajkapooret *et al.* (2008)¹⁴ and Gillette (2000),¹⁵ herbal remedies are becoming increasingly popular as complementary medications for the treatment and prevention of a wide range of illnesses, most of which are caused by exposure to stress or pollutants. According to research done by Subramonium and Puspangadan (1999)¹⁶ and Hatem *et al.* (201) a number of plant components have the ability to reduce the liver damage brought on by various chemical agents. It has also been demonstrated in several experiments that *A. conyzoides*' various sections cannot be extracted using aqueous or alcoholic methods. The liver's membrane stability and ability to operate is preserved by extracts. These biochemical restoration by plant extracts may be caused by cytochrome P-450 inhibition, promotion of its glucuronidation of toxic metabolites, or directly scavenging activity of reactive intermediate molecules responsible for the hepatocytes insert. Yellow pigment bilirubin (unconjugated) is created when heme is catabolized. Prior to being secreted into bile, hepatocytes conjugate bilirubin with glucuronic acid (conjugated bilirubin) to make it watersoluble and so readily excretable. Hepatocytes' failure to perform the conjugation process under exposure to hepatotoxins led to an increase in the levels of unconjugated bilirubin. The pretreatment group that had been given acetone and n-hexane extracts of *A. conyzoides* had considerably lower levels of these enzymes than the group that had been exposed to APAP, demonstrating that the extracts continue to maintain the liver's ability to operate. Another well-known very effective hepatoprotective compound is silymarin, a standardised extract of *Silybum marianum* (Compositae). The liver damage caused by carbon tetra chloride, D-galN, and APAP in rat models was averted by this compound, which reverses hepatotoxin-induced abnormalities in biochemical parameters (Ramellini and Meldoles, 1976;¹⁸ Bahati *et al.*, 2006)¹⁹. This compound has thus far undergone the most extensive investigation of all plant compounds. Similar findings on acetaminophen-induced liver damage were previously reported by Hattori *et al.* (1990)²⁰ and Emmanuel *et al.* (1995)²¹. The elevated levels of BUN and CR in the APAP-treated group might be attributed to protein metabolism. Albumin, transferrin, and transthyretin are examples of the negative acute-phase proteins, whose production and secretion are lowered (Trey and Kushner, 1995)²². According to the study's findings, groups of rats administered APAP alone had lower levels of albumin and total protein than control groups. Administration of *Ageratum conyzoides* extracts, however, reduced the impact, reestablish the levels of SGPT, SGOT, LDH, and bilirubin as a sign that the plasma membrane has stabilised and that the hepatic tissue damages brought on by APAP have been repaired.

IV. CONCLUSION

Our study indicated that the alcoholic extract of *Ageratum conyzoides* isolated compound could be beneficial as hepatoprotective agents and showed significant result. These results suggest that the application of *Ageratum conyzoides* extract in CCl₄ induced toxicity can be reversed. However extensive research is required to explore for safer use of herbal preparation.

V. REFERENCES

- [1] Adebayo AH, Zeng GZ, Zhang YM, Ji CJ, Afolabi AA, Tan NH. (2010) Toxicological evaluation of precocene II isolated from *Ageratum conyzoides* L. (Asteraceae) in Sprague Dawley rats. African Journal of Biotechnology.; 9(20):2938-44.

- [2] Akilavalli N, Radhika J, Brindha P.(2011)Hepato protective activity of Ocimum sanctum Linn. Against lead induced toxicity in abino rats. Asian J.Pharm. Clin.Res.;4:84-87
- [3] Bahati JR, Goyal RK, Shah GB. (2006)Hepatoprotective activity of Hemidesmus indicus R.Br in rats. Ind J Exp Biol.; 44: 399-402.
- [4] Emmanuel J. Kambisi L, Afolayan A.J.(2000) An Ethnobotanical Study of Plants Use for the treatment of STD's. Guruye District. Zimbabwe, 1995. Gillette JR. Laboratory of Chemical Pharmacology, National Heart, Lung and Blood Institute, NIH: a short history. Annu Rev Pharmacol Toxicol.; 40:18-41.
- [5] Gonzalez AG, Aguiar ZE, Grillo TA, Luis JG, Rivera A, Calle J.(1991) Chromenes from Ageratum conyzoides. Phytochemistry.; 30: 1137-9.
- [6] H. Oxidative Stress, Wolf. PC (1999) Biochemical of liver diseases India J Clin. Biochem.;14:59-90
- [7] Handa SS, Sharma A, Chakraborty KK. (1989);Natural products and plants as liver protecting drugs. Fitoterapia. 57:307-351
- [8] Hatem S. Farghaly and Mohammed A. Hussein.(2010)Protective Effect of Curcumin Against Paracetamol-induced Liver Damage. Australian Journal of Basic and Applied Sciences.; 4(9): 4266-4274
- [9] Hattori S.M., Makkai H.P.S, Becker K. (1990). Feed Value of Some Browse plants. Central Zone of Delta State, Nigeria.; 39:97-104.
- [10] Jander EA, Machado KCCA,(2008) Evolutionary ecology of figs and their associates: Recent progress and outstanding Puzzles.Ann Rev Evol.Syst.; 39:439-458
- [11] K., A. Mansouri, G. Bosku, P. Kefalas. (2005)Tea and herbal infusions: their antioxidant activity and phenolic profile. Food Chemistry.; 89:27-36.
- [12] KaplowitzN,(1996) Tsukamoto and liver disease. Progress in Liver Diseases.; 14:131-159
- [13] Manik Sharma and Abid Rashid (2017)Protective Effect of Leaves of Ficus carica Against Carbon Tetrachloride-Induced Hepatic Damage in Rats.UK 2017Journal of Pharmaceutical and Biosciences - Vol.5(1),06-11,
- [14] Mohan GK, Pallavi E, Ravi Kumar B, Ramesh M, Venkatesh S.(2007) Hepatoprotective activity of Ficus carica Linn leaf extract against carbon tetrachloride-induced hepatotoxicity in rats. Daru. ;15:162-166
- [15] Premitha Abraham P, Wilted G, Ramakrishna B.(1999) Decreased activity of hepatic alkaline proprieties in rats with carbon tetrachloride induced liver cirrhosis. Indian. J Exp. Biol.; 37:1234-1244
- [16] Rose JA, Kasum CM. (2002)Dietary flavonoids: bioavailability metabolic effects, and safety. Annu Rev Nutr; 22:19-34.
- [17] Saleem TSM, Chetty CM, Ramkanath S, Rajan VST, Kumar KM, GauthmanK. (2010)Hepatoprotective herbs-a review. International Journal of Research in pharmaceutical Science; 1; 1 1-5.
- [18] Tibrell J.(2001)Introduction to toxicology. 3rd ed. Taylor and Francis. New York.;57-71
- [19] Vaya J, Mahmood S.2006 Flavonoid content in leaf extracts of the fig (Ficus carica L.) carob (Ceratonia siliqua L.) and Pistachio (Pistacia lentiscus L. Biofactors.;22:19-34
- [20] Ita SO, Akpanyung EO, Umoh BI, Ben EE, Ukafia SO.(2009); Acetaminophen induced hepatic toxicity: Protective Role of Ageratum conyzoides. Pakistan Journal of Nutrition. 8(7):928-32.
- [21] Ita, S.O., Etim, O.E., Ben, E.E., Ekpo, O.F. (2007); Haemato-poietic properties of ethanolic leaf extract of Ageratum conyzoides (Goat weed) in albino rats. Nigerian Journal of Physiology and Science, 22: 83-87.
- [22] Jagetia GC, Shirwaikar A, Rao SK, Bhilegaonkar PM. (2003) Evaluation of the radio-protective effect of Ageratum conyzoides L. extract in mice, exposed to different doses of gamma radiation. J. Pharm. Pharmacol.; 55(8):1151-8.
- [23] John AA, Bamidele FP, Oloyede AO, Seun OAO. (2011) Some protective effects of aqueous leaf extract of Azadirachta indica on paracetamol-induced hepatotoxicity in adult wistar rats. Am. J. of Trop. Med. & Public Health.; 1(3):97-106.
- [24] Kim NK, Vasminah WG, Frejar EF, Goldman AI, Theologides A.(1977) Value of alkaline phosphatase, 5'-nucleotidase, gamma-glutamyl transferase and glutamate dehydrogenase activity measurements (Single and Combined) is serum in diagnosis of metastases to the liver. Clinical Chem.; 23:2034-2038.